

First edition
2017-03

AMENDMENT 1
2021-03

**Microbiology of the food chain —
Horizontal method for determination
of hepatitis A virus and norovirus
using real-time RT-PCR —**

**Part 1:
Method for quantification**

AMENDMENT 1

*Microbiologie de la chaîne alimentaire — Méthode horizontale pour
la recherche des virus de l'hépatite A et norovirus par la technique
RT-PCR en temps réel —*

Partie 1: Méthode de quantification

AMENDEMENT 1



Reference number
ISO 15216-1:2017/Amd.1:2021(E)

© ISO 2021



COPYRIGHT PROTECTED DOCUMENT

© ISO 2021

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
CP 401 • Ch. de Blandonnet 8
CH-1214 Vernier, Geneva
Phone: +41 22 749 01 11
Email: copyright@iso.org
Website: www.iso.org

Published in Switzerland

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation of the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT), see www.iso.org/iso/foreword.html.

This document was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 9, *Microbiology*, in collaboration with the European Committee for Standardization (CEN) Technical Committee CEN/TC 463, *Microbiology of the food chain*, in accordance with the Agreement on technical cooperation between ISO and CEN (Vienna Agreement).

A list of all parts in the ISO 15216 series can be found on the ISO website.

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at www.iso.org/members.html.

STANDARDSISO.COM : Click to view the full PDF of ISO 15216-1:2017/Amd 1:2021

Microbiology of the food chain — Horizontal method for determination of hepatitis A virus and norovirus using real-time RT-PCR —

Part 1: Method for quantification

AMENDMENT 1

5.2.16

Replace the text with the following:

Ethylenediaminetetraacetic acid (EDTA) disodium dihydrate.

8.2.5, first paragraph

Replace the text with the following:

This document is appropriate for water bottles with volumes up to 2 l. The entire contents of the bottle should be tested. For each sample, record the volume tested.

B.2.2

Replace the text with the following:

Mix the components together.

Store at room temperature in a dark glass bottle for a maximum of 12 months.

B.7.1

Replace the text with the following:

Ethylenediaminetetraacetic acid (EDTA) disodium dihydrate	(18,6 ± 0,2) g
---	----------------

Water (5.2.1)	as required
---------------	-------------

B.7.2

Replace the first sentence with the following:

Dissolve the EDTA disodium dihydrate in (90 ± 1) ml water.

E.1, first paragraph

Replace the second sentence with the following:

Mengo virus strain MC₀ (CECT 100 000)² is a recombinant (deletant) virus which lacks the poly(C) tract in comparison to the wild-type mengo virus, with identical growth properties to those of the wild-type virus but with an avirulent phenotype.

E.1, Footnote 2

Replace the first sentence with the following:

CECT 100 000 and ATCC® CCL-2™ are trademarks of products supplied by the Spanish Type Culture Collection and American Type Culture Collection respectively.

F.1.2, Footnote 3

Replace the text with the following:

BioMerieux NucliSens® is the trade name of a product supplied by bioMerieux. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of the product named. Equivalent products can be used if they can be shown to lead to the same results.

F.2

Replace the four subclauses with the following subclauses:

F.2.1 Magnetic rack for 1,5 ml tubes.

F.2.2 Thermoshaker or equivalent apparatus for shaking 1,5 ml tubes at $(60 \pm 2) ^\circ\text{C}$ and approximately 1 400 oscillations min⁻¹.

F.3

Replace the text with the following:

Add $(2 \pm 0,1)$ ml of NucliSens® lysis buffer to a tube. Add (500 ± 10) µl of sample (BMS) or entire sample (other matrices) and mix by vortexing briefly.

Incubate for (10 ± 1) min at room temperature.

Add $(50 \pm 2,5)$ µl of well-mixed magnetic silica solution to the tube and mix by vortexing briefly.

Incubate for (10 ± 1) min at room temperature.

Centrifuge for (120 ± 10) s at 1 500g or allow silica to sediment using a magnetic rack then carefully discard supernatant by, for example, aspiration.

Add (400 ± 10) µl wash buffer 1 and resuspend the pellet by pipetting or vortexing, taking care to avoid foaming.

Transfer suspension to a clean 1,5 ml tube. Cap tube and wash silica for (30 ± 2) s by vortexing. After washing, allow silica to sediment using the magnetic rack. Discard supernatant by, for example, aspiration.