
**Oil of ylang-ylang [*Cananga odorata*
(Lam.) Hook. f. et Thomson forma
genuina]**

*Huile essentielle d'ylang-ylang [Cananga odorata (Lam.) Hook. f. et
Thomson forma genuina]*

STANDARDSISO.COM : Click to view the full PDF of ISO 3063:2004



PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

STANDARDSISO.COM : Click to view the full PDF of ISO 3063:2004

© ISO 2004

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

Published in Switzerland

Contents

Page

Foreword.....	iv
1 Scope.....	1
2 Normative references	1
3 Terms and definitions	1
4 Requirements	1
4.1 Appearance.....	1
4.2 Colour.....	1
4.3 Odour.....	1
4.4 Physical and chemical requirements	1
4.5 Chromatographic profile	1
4.6 Flashpoint	2
5 Sampling	2
6 Test methods	2
6.1 Relative density at 20 °C d_{20}^{20}	2
6.2 Refractive index at 20 °C	2
6.3 Optical rotation at 20 °C	2
6.4 Acid value	2
6.5 Ester value	2
6.6 Chromatographic profile	2
7 Packaging, labelling, marking and storage	2
Annex A (informative) Typical chromatograms of the analysis by gas chromatography of oil of ylang-ylang [<i>Cananga odorata</i> (Lam.) Hook. f. et Thomson forma <i>genuina</i>]	6
Annex B (informative) Flashpoint.....	11
Bibliography	12

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 3063 was prepared by Technical Committee ISO/TC 54, *Essential oils*.

This second edition cancels and replaces the first edition (ISO 3063:1983), which has been technically revised.

STANDARDSISO.COM : Click to view the full PDF of ISO 3063:2004

Oil of ylang-ylang [*Cananga odorata* (Lam.) Hook. f. et Thomson forma *genuina*]

1 Scope

This International Standard specifies certain characteristics of the oil of ylang-ylang [*Cananga odorata* (Lam.) Hook. f. et Thomson forma *genuina*] from Madagascar, Mayotte and Comores, in order to facilitate assessment of its quality.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO/TR 210, *Essential oils — General rules for packaging, conditioning and storage*

ISO/TR 211, *Essential oils — General rules for labelling and marking of containers*

ISO 212, *Essential oils — Sampling*

ISO 279, *Essential oils — Determination of relative density at 20 °C — Reference method*

ISO 280, *Essential oils — Determination of refractive index*

ISO 592, *Essential oils — Determination of optical rotation*

ISO 709, *Essential oils — Determination of ester value*

ISO 1242, *Essential oils — Determination of acid value*

ISO 11024-1, *Essential oils — General guidance on chromatographic profiles — Part 1: Preparation of chromatographic profiles for presentation in standards*

ISO 11024-2, *Essential oils — General guidance on chromatographic profiles — Part 2: Utilization of chromatographic profiles of samples of essential oils*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

3.1

oil of ylang-ylang

essential oil obtained by steam distillation of the fresh flowers of *Cananga odorata* (Lam.) Hook. f. et Thomson forma *genuina*, of the Annonaceae family, growing mainly in Madagascar, Mayotte and Comores

NOTE 1 This volatile product is not generally collected as a whole oil, but in five successive fractions during the course of distillation. These five fractions, known respectively as "Extra super", "Extra", "First", "Second" and "Third", are the oils usually found in the trade.

NOTE 2 For information on the CAS number, see ISO/TR 21092.

4 Requirements

4.1 Appearance

Liquid.

4.2 Colour

Pale yellow to dark yellow.

4.3 Odour

Characteristic, floral and recalling jasmine.

4.4 Physical and chemical requirements

See Table 1.

4.5 Chromatographic profile

Analysis of the essential oil shall be carried out by gas chromatography. In the chromatogram obtained, the representative and characteristic components shown in Table 2 shall be identified. The proportions

of these components, indicated by the integrator, shall be as shown in Table 2. This constitutes the chromatographic profile of the essential oil.

4.6 Flashpoint

Information on the flashpoint is given in Annex B.

5 Sampling

See ISO 212.

Minimum volume of test sample: 25 ml

NOTE This volume allows each of the tests specified in this International Standard to be carried out at least once.

6 Test methods

6.1 Relative density at 20 °C d_{20}^{20}

See ISO 279.

6.2 Refractive index at 20 °C

See ISO 280.

6.3 Optical rotation at 20 °C

See ISO 592.

6.4 Acid value

See ISO 1242.

6.5 Ester value

See ISO 709.

6.6 Chromatographic profile

See ISO 11024-1 and ISO 11024-2.

7 Packaging, labelling, marking and storage

See ISO/TR 210 and ISO/TR 211.

Table 1 — Physical and chemical requirements

Characteristics	Fractions								
	Extra super	Extra		First		Second		Third	
	Comores and Mayotte	Comores and Mayotte	Mada- gascar	Comores and Mayotte	Mada- gascar	Comores and Mayotte	Mada- gascar	Comores and Mayotte	Mada- gascar
Relative density at 20 °C d_{20}^{20}									
Min.	0,970	0,955	0,950	0,938	0,933	0,925	0,922	0,906	0,906
Max.	0,990	0,976	0,965	0,960	0,949	0,945	0,942	0,925	0,925
Refractive index at 20 °C									
Min.	1,497	1,498	1,493	1,501	1,495	1,502	1,496	1,503	1,502
Max.	1,505	1,506	1,509	1,509	1,510	1,511	1,511	1,513	1,513
Optical rotation at 20 °C									
Min.	– 33°	– 40°	– 42°	– 46°	– 46°	– 60°	– 58°	– 72°	– 70°
Max.	– 12,5°	– 20°	– 20°	– 25°	– 24°	– 35°	– 30°	– 45°	– 45°
Acid value	< 2	< 2	< 2	< 2	< 2	< 2	< 2	< 2	< 2
Ester value									
Min.	160	140	125	100	90	75	65	45	40
Max.	200	185	160	160	125	115	95	75	70

Table 2 — Chromatographic profile

Component	Fractions								
	Extra super	Extra		First		Second		Third	
	Comores and Mayotte	Comores and Mayotte	Mada- gascar	Comores and Mayotte	Mada- gascar	Comores and Mayotte	Mada- gascar	Comores and Mayotte	Mada- gascar
Prenyl acetate									
Min.	1,5	1,0	0,6	0,3	0,2	0,2	0,1	0,1	traces
Max.	3,2	2,3	2,2	1,8	1,0	0,9	0,5	0,2	0,2
p-Cresyl methyl ether									
Min.	7,0	5,0	7,0	3,0	5,0	2,0	1,0	0,1	0,1
Max.	13,0	13,0	16,0	8,5	10,0	5,0	4,6	1,0	1,4
Methyl benzoate									
Min.	4,5	4,0	4,5	1,5	3,0	1,0	1,0	0,1	0,1
Max.	8,0	6,5	9,0	5,5	5,0	3,5	3,0	0,8	0,9
Linalool									
Min.	8,0	7,0	15,0	3,0	12,0	2,0	4,0	0,1	0,6
Max.	13,0	12,0	24,0	10,0	19,0	6,0	9,5	2,0	4,0
Benzyl acetate									
Min.	14,0	11,0	5,5	6,0	2,8	4,0	0,5	0,5	0,1
Max.	20,0	17,5	14,0	14,0	10,0	8,8	5,0	3,0	2,2
Geraniol									
Min.	0,1	0,1	1,3	0,1	1,6	0,1	0,7	traces	0,2
Max.	0,7	0,5	3,0	0,3	2,6	0,3	2,4	0,1	0,8
Geranyl acetate									
Min.	2,0	2,5	7,0	2,0	8,0	1,7	5,6	0,4	1,0
Max.	6,0	6,0	14,0	5,0	15,0	6,0	12,0	3,0	6,6
E-Cinnamyl acetate									
Min.	4,0	3,0	0,5	2,2	0,5	2,0	0,4	0,5	0,1
Max.	6,0	6,5	3,0	5,0	2,0	4,8	2,2	2,5	2,0
β-Caryophyllene									
Min.	2,0	2,5	2,5	4,0	5,5	4,8	10,0	5,0	12,0
Max.	6,0	8,0	8,5	10,0	12,0	14,0	17,0	15,0	19,0
D-Germacrene									
Min.	9,0	14,0	5,0	10,0	9,5	16,0	13,0	20,0	15,0
Max.	15,0	20,0	15,0	24,0	18,0	28,0	28,0	35,0	34,0
(E,E)-α-Farnesene									
Min.	2,0	6,5	1,0	7,0	3,0	14,0	5,0	12,0	9,0
Max.	6,0	15,0	5,0	18,0	8,0	21,0	11,5	29,0	25,0

Table 2 (continued)

Component	Fractions								
	Extra super	Extra		First		Second		Third	
	Comores and Mayotte	Comores and Mayotte	Mada- gascar	Comores and Mayotte	Mada- gascar	Comores and Mayotte	Mada- gascar	Comores and Mayotte	Mada- gascar
(E,E)-Farnesol									
Min.	0,8	0,8	0,5	0,8	0,1	0,8	1,2	0,8	1,2
Max.	1,5	1,6	3,0	2,0	2,5	3,0	3,5	3,0	4,0
Benzyl benzoate									
Min.	3,0	4,0	3,5	4,2	4,5	4,5	6,0	4,0	4,8
Max.	6,0	6,0	8,0	9,2	8,0	7,8	10,0	8,0	8,5
(E,E)- Farnesyl acetate									
Min.	1,0	1,0	0,5	1,0	1,0	1,0	1,2	1,5	1,7
Max.	3,0	3,0	3,0	4,0	2,0	3,5	3,5	5,0	5,0
Benzyl salicylate									
Min.	1,5	2,0	1,2	2,0	1,6	2,0	1,8	2,5	2,0
Max.	3,5	3,8	4,0	4,0	4,0	4,0	4,0	4,8	5,0

Annex A

(informative)

Typical chromatograms of the analysis by gas chromatography of oil of ylang-ylang [*Cananga odorata* (Lam.) Hook. f. et Thomson forma *genuina*]





Figure A.1 — Typical chromatogram of oil of ylang-ylang Extra Comores taken on an apolar column


**Peak identification**

- | | |
|---------------------------------|--|
| 1 Prenyl acetate | 9 β -Caryophyllene |
| 2 <i>p</i> -Cresyl methyl ether | 10 D-Germacrene |
| 3 Methyl benzoate | 11 (<i>E,E</i>)- α -Farnesene |
| 4 Linalool | 12 (<i>E,E</i>)-Farnesol |
| 5 Benzyl acetate | 13 Benzyl benzoate |
| 6 Geraniol | 14 (<i>E,E</i>)-Farnesyl acetate |
| 7 Geranyl acetate | 15 Benzyl salicylate |
| 8 <i>E</i> -Cinnamyl acetate | |

Operating conditions

Column: silica capillary, length 20 m, internal diameter 0,1 mm
 Stationary phase: methyl siloxane
 Film thickness: 0,40 μ m
 Oven temperature: isothermal at 80 °C for 2 min, then temperature programming from 80 °C to 170 °C at a rate of 4 °C/min, and from 170 °C to 305 °C at a rate of 15 °C/min, and isothermal at 305 °C for 2 min
 Injector temperature: 260 °C
 Detector temperature: 300 °C
 Detector: flame ionization type
 Carrier gas: hydrogen
 Volume injected: 0,2 μ l
 Carrier gas flow rate: 0,6 ml/min
 Split ratio: 1/120

Figure A.2 — Typical chromatogram of oil of ylang-ylang “Third” Comores taken on an apolar column


**Peak identification**

- | | |
|---------------------------------|--|
| 1 Prenyl acetate | 9 β -Caryophyllene |
| 2 <i>p</i> -Cresyl methyl ether | 10 δ -Germacrene |
| 3 Methyl benzoate | 11 (<i>E,E</i>)- α -Farnesene |
| 4 Linalool | 12 (<i>E,E</i>)-Farnesol |
| 5 Benzyl acetate | 13 Benzyl benzoate |
| 6 Geraniol | 14 (<i>E,E</i>)-Farnesyl acetate |
| 7 Geranyl acetate | 15 Benzyl salicylate |
| 8 <i>E</i> -Cinnamyl acetate | 16 Geranyl benzoate |

Operating conditions

Column: capillary, length 50 m, internal diameter 0,2 mm
 Stationary phase: poly (dimethyl siloxane)
 Film thickness: 0,25 μ m
 Oven temperature: programme of temperature from 65 °C to 230 °C at a rate of 2 °C/min
 Injector temperature: 230 °C
 Detector temperature: 250 °C
 Detector: flame ionization type
 Carrier gas: hydrogen
 Volume injected: 0,2 μ l
 Carrier gas flow rate: 1,1 ml/min
 Split ratio: 1/100

Figure A.3 — Typical chromatogram of oil of ylang-ylang “Third” Madagascar taken on an apolar column


**Peak identification**

- | | |
|---------------------------------|---------------------------------------|
| 1 Prenyl acetate | 9 (<i>E,E</i>)- α -Farnesene |
| 2 <i>p</i> -Cresyl methyl ether | 10 Geraniol |
| 3 Linalool | 11 <i>E</i> -Cinnamyl acetate |
| 4 Methyl benzoate | 12 (<i>E,E</i>)-Farnesyl acetate |
| 5 β -Caryophyllene | 13 (<i>E,E</i>)-Farnesol |
| 6 Benzyl acetate | 14 Benzyl benzoate |
| 7 δ -Germacrene | 15 Benzyl salicylate |
| 8 Geranyl acetate | |

Operating conditions

Column: silica, capillary, length 20 m, internal diameter 0,1 mm
 Stationary phase: bonded polyethylene glycol
 Film thickness: 0,20 μ m
 Oven temperature: isothermal at 65 °C for 1 min, then temperature programming from 65 °C to 190 °C at a rate of 10 °C/min, and isothermal 190 °C for 2 min, and from 190 °C to 240 °C, at a rate of 20 °C/min, and isothermal 240 °C for 6 min
 Injector temperature: 250 °C
 Detector temperature: 280 °C
 Detector: flame ionization type
 Carrier gas: hydrogen
 Volume injected: 0,2 μ l
 Carrier gas flow rate: 0,4 ml/min
 Split ratio: 1/150

Figure A.4 — Typical chromatogram of oil of ylang-ylang Extra Comores taken on a polar column

**Peak identification**

- | | |
|---------------------------------|---------------------------------------|
| 1 Prenyl acetate | 9 (<i>E,E</i>)- α -Farnesene |
| 2 <i>p</i> -Cresyl methyl ether | 10 Geraniol |
| 3 Linalool | 11 <i>E</i> -Cinnamyl acetate |
| 4 Methyl benzoate | 12 (<i>E,E</i>)-Farnesyl acetate |
| 5 β -Caryophyllene | 13 (<i>E,E</i>)-Farnesol |
| 6 Benzyl acetate | 14 Benzyl benzoate |
| 7 δ -Germacrene | 15 Benzyl salicylate |
| 8 Geranyl acetate | |

Operating conditions

Column: silica, capillary, length 20 m, internal diameter 0,1 mm
 Stationary phase: bonded polyethylene glycol
 Film thickness: 0,20 μ m
 Oven temperature: isothermal at 65 °C for 1 min, then temperature programming from 65 °C to 190 °C at a rate of 10 °C/min, and isothermal 190 °C for 2 min, and from 190 °C to 240 °C, at a rate of 20 °C/min, and isothermal 240 °C for 6 min
 Injector temperature: 250 °C
 Detector temperature: 280 °C
 Detector: flame ionization type
 Carrier gas: hydrogen
 Volume injected: 0,2 μ l
 Carrier gas flow rate: 0,4 ml/min
 Split ratio: 1/150

Figure A.5 — Typical chromatogram of oil of ylang-ylang “Third” Comores taken on a polar column